

AD-A177 292


ON A LIMIT THEOREM AND INVARIANCE PRINCIPLE FOR
SYMMETRIC STATISTICS (U) NORTH CAROLINA UNIV AT CHAPEL
HILL CENTER FOR STOCHASTIC PROC V MANDREKAR JUL 86
TR-142 AFOSR-TR-87-0121 F49620-85-C-0144 F/G 12/1

1/1

UNCLASSIFIED

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

(2)

AD-A177 292

REPORT DOCUMENTATION PAGE				
1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED		1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY NA		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for Public Release; Distribution Unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE NA		5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR- 87-0121		
1. PERFORMING ORGANIZATION REPORT NUMBER Technical Report No. 142		7a. NAME OF MONITORING ORGANIZATION AFOSR/NM		
1a. NAME OF PERFORMING ORGANIZATION University of North Carolina		8b. OFFICE SYMBOL (If applicable)		7b. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB, DC 20332-6448
1c. ADDRESS (City, State and ZIP Code) Center for Stochastic Processes, Statistics Department, Phillips Hall 039-A, Chapel Hill, NC 27514		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F49620 85 C 0144		
1a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR		8b. OFFICE SYMBOL (If applicable)		10. SOURCE OF FUNDING NOS.
8c. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB, DC		PROGRAM ELEMENT NO. 6.1102F		PROJECT NO. 2304
11. TITLE (Include Security Classification) "On a limit theorem and invariance principle for symmetric statistics"		TASK NO. HS		WORK UNIT NO.
12. PERSONAL AUTHOR(S) Mandrekar, V.				
13a. TYPE OF REPORT technical		13b. TIME COVERED FROM 9/85 TO 8/86		14. DATE OF REPORT (Yr., Mo., Day) July 1986
15. PAGE COUNT 6				
16. SUPPLEMENTARY NOTATION				
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB. GR.		
XXXXXXXXXXXXXXXXXXXX		Keywords:		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) The purpose of this note is to give a direct proof of some recent important results of E.B. Dynkin and A. Mandelbaum [2]. This also provides immediately the results in [3] with a very simple proof. This is achieved by avoiding the use of Poisson process. Let us set up some notation. Let (L, Σ, μ) be a probability space and (X^k, Σ^k, μ^k) be the k -fold product probability space. Let $h_k(x_1, \dots, x_k)$ be a symmetric function of k -variables. We call it canonical if $\int h_k(x_1, \dots, x_{k-1}, y) d\mu = 0$ for all $x_1, \dots, x_{k-1} \in X^{k-1}$. Let X_1, \dots, X_n be a i.i.d. X -valued random variable on a probability space with distribution μ .				
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL Daggy Ravitch Maj Crawley		22b. TELEPHONE NUMBER (Include Area Code) 919-962-2367 7673 8023		22c. OFFICE SYMBOL AFOSR/NM

DD FORM 1473, 83 APR

EDITION OF 1 JAN 73 IS OBSOLETE.


UNCLASSIFIED

772 FILE COPY

CENTER FOR STOCHASTIC PROCESSES

AFOSR-TR- 87-0121

Department of Statistics
University of North Carolina
Chapel Hill, North Carolina



ON A LIMIT THEOREM AND INVARIANCE
PRINCIPLE FOR SYMMETRIC STATISTICS

BY

V. Mandrekar

Approved for public release;
distribution unlimited.

Technical Report No. 142

July 1986

87 2 20 1986

ON A LIMIT THEOREM AND INVARIANCE PRINCIPLE FOR SYMMETRIC STATISTICS

by

V. Mandrekar*

Department of Statistics and Probability
Michigan State University
East Lansing, MI 68824

and

Center for Stochastic Processes
Department of Statistics
University of North Carolina
Chapel Hill, NC 27514

Accession For	
NTIS CRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	



*This research supported by ONR N00014-85-K-0150 and the Air Force Office of Scientific Research Contract No. F49620 85C 0144.

0. Introduction: The purpose of this note is to give a direct proof of some recent important results of E.B. Dynkin and A. Mandelbaum [2]. This also provides immediately the results in [3] with a very simple proof. This is achieved by avoiding the use of Poisson process. Let us set up some notation. Let (X, Σ, μ) be a probability space and (X^k, Σ^k, μ^k) be the k -fold product probability space. Let $h_k(x_1, \dots, x_k)$ be a symmetric function of k -variables. We call it canonical if $\int h_k(x_1, \dots, x_{k-1}, y) d\mu = 0$ for all $x_1, \dots, x_{k-1} \in X^{k-1}$. Let X_1, \dots, X_n be a i.i.d. X -valued random variable on a probability space with distribution μ . As in [2], define

$$\sigma_k^n(h_k) = \sum_{1 \leq s_1 < \dots < s_k \leq n} h_k(X_{s_1}, \dots, X_{s_k}), \text{ for } k \leq n$$

$$= 0 \quad \text{for } k > n.$$

Let $H = \{(h_0, h_1, \dots) : h_k \text{ canonical and } \sum_{k=1}^{\infty} \frac{1}{k!} \|h_k\|_2^2 < \infty\}$ where h_0 is a constant and $\|\cdot\|_2$ is the norm in $L^2(X^k, \Sigma^k, \mu^k)$. On H define

$\|h\|^2 = \sum_{k=0}^{\infty} \|h_k\|_2^2 / k!$. H is the so-called exponential (Fock) space of $L_0^2(X, \Sigma, \mu)$ ($\phi \in L^2(X, \Sigma, \mu)$ with $E\phi(X) = 0$). It is a Hilbert space under coordinate addition, scalar multiplication and $\|\cdot\|$. For each $\phi \in L_0^2(X, \Sigma, \mu)$, $h^\phi \in H$ with $h_k^\phi = \phi(x_1), \dots, \phi(x_k)$. It can be easily seen that $\text{sp}\{h^\phi : \phi \in L_0^2(X, \Sigma, \mu)\}$ is dense in H . Define for each $h \in H$,

$$(0.1) \quad Y_n(h) = \sum_{k=0}^{\infty} n^{-k/2} \sigma_k^n(h_k).$$

Since $\sigma_k^n(h_k) = 0$ for $k > n$, this is a finite sum. Also, let

$$(0.2) \quad Y_n^t(h) = \sum_{k=0}^{\infty} n^{-k/2} \sigma_k^{[nt]}(h_k).$$

The main purpose is to show directly that $Y_n(h) \xrightarrow{\mathcal{D}} \sum_{k=0}^{\infty} \frac{I_k(h_k)}{k!}$ where $\xrightarrow{\mathcal{D}}$ denotes convergence in distribution and $I_k(h_k)$ denotes Ito-Wiener multiple

integral of h_k with respect to Gaussian random measure W with $EW(A)W(A') = \mu(A \cap A')$.

In the next section we discuss the convergence of $Y_n^t(h)$. We observe that for $\phi \in L_0^2(X, \Sigma, \mu)$

$$\begin{aligned} Y_n(h^\phi) &= \sum_{k=0}^n n^{-k/2} \sum_{1 \leq s_1 < \dots < s_k \leq n} \phi(X_{s_1}) \dots \phi(X_{s_k}) \\ &= \sum_{k=0}^n \sum_{1 \leq s_1 < \dots < s_k \leq n} \frac{\phi(X_{s_1})}{\sqrt{n}} \dots \frac{\phi(X_{s_k})}{\sqrt{n}} \\ &= \prod_{i=1}^n \left(1 + \frac{\phi(X_i)}{\sqrt{n}}\right). \end{aligned}$$

Let us observe that for any $\varepsilon > 0$,

$$\sum_j P(|\phi(X_j)| > \sqrt{\varepsilon j}) = \sum_j P(|\phi(X_1)|^2 > \varepsilon j) \leq \|\phi\|_2^2 < \infty.$$

Hence by Borel-Cantelli lemma, a.s. (for $j \leq n$)

$$|\phi(X_j)| \leq \sqrt{\varepsilon j} \leq \sqrt{\varepsilon} \sqrt{n} \quad \text{for } j \geq \text{some } N(\omega) \quad (N(\omega) < \infty).$$

$$\text{But } \prod_{i=1}^n \left(1 + \frac{\phi(X_i)}{\sqrt{n}}\right) = \prod_{i=1}^{N(\omega)} \left(1 + \frac{\phi(X_i)}{\sqrt{n}}\right) \prod_{i=N(\omega)+1}^n \left(1 + \frac{\phi(X_i)}{\sqrt{n}}\right) \quad \text{giving for a.s. } \omega, \text{ so}$$

$$\lim_n Y_n(h^\phi) = \lim_n \prod_{i=N(\omega)}^n \left(1 + \frac{\phi(X_i)}{\sqrt{n}}\right). \quad \text{Thus WLOG, we can assume for } n \text{ large}$$

$$\left|\frac{\phi(X_j)}{\sqrt{n}}\right| < 1 \quad \text{a.s. for all } j \leq n \text{ and } Y_n(h^\phi) = \prod_{i=1}^n \left(1 + \frac{\phi(X_i)}{\sqrt{n}}\right). \quad \text{Taking log on both sides and expanding } \log(1+x) \text{ we have}$$

$$\log \prod_{i=1}^n \left(1 + \frac{\phi(X_i)}{\sqrt{n}}\right) = \sum_{i=1}^n \frac{\phi(X_i)}{\sqrt{n}} - \frac{1}{2} \sum_{i=1}^n \frac{\phi(X_i)^2}{n} + \varepsilon_n(\phi)$$

where $\varepsilon_n(\phi) \xrightarrow{P} 0$ by the WLLN and since $\max_i \left|\frac{\phi(X_i)}{\sqrt{n}}\right| \xrightarrow{P} 0$ by Chebychev's Inequality,

i.e. the $(Y_n(h^\phi)) \xrightarrow{D} \exp[I_1(\phi) - \frac{1}{2}\|\phi\|_2^2]$. Using Cramér-Wold device and the above argument we get

0.3 Lemma: For any finite subset $\{\phi_1, \dots, \phi_k\} \subseteq L^2(X, \Sigma, \mu)$

$$(Y_n(h^{\phi_1}), \dots, Y_n(h^{\phi_k})) \xrightarrow{D} (\exp(I_1(\phi_1) - \frac{1}{2}\|\phi_1\|_2^2), \dots, \exp(I_1(\phi_k) - \frac{1}{2}\|\phi_k\|_2^2)).$$

As a consequence, we get for $\{\phi_i, i \in I\}$ a finite subset of $L^2(X, \Sigma, \mu)$ and

$$\{c_i, i \in I\} \subseteq \mathbb{R},$$

$$(0.3)' \quad Y_n\left(\sum_{i \in I} c_i h^{\phi_i}\right) \xrightarrow{D} \sum_{k=0}^{\infty} \frac{I_k\left(\sum_{i \in I} c_i h^{\phi_i}\right)_k}{k!}.$$

We now observe that for $h, h' \in H$,

$$(0.4) \quad E[Y_n(h) - Y_n(h')]^2 = \sum_k \binom{n}{k} n^{-k} \|h_k - h'_k\|^2 \leq E\|h - h'\|^2,$$

since $E\sigma_k^n(h_k - h'_k)\sigma_\ell^n(h_\ell - h'_\ell) = \binom{n}{k} \|h_k - h'_k\|^2 \delta_{k\ell}$ by ([2], p. 744). Also,

$$(0.5) \quad E\left(\sum_{k=0}^{\infty} I_k(h_k)/k! - \sum_{k=0}^{\infty} \frac{I_k(h'_k)}{k!}\right)^2 = \|h - h'\|^2.$$

Thus we get

(0.6) Theorem: For any $h \in H$,

$$Y_n(h) \xrightarrow{D} W(h) = \sum_{k=0}^{\infty} \frac{I_k(h_k)}{k!}$$

Proof: Let $h \in H$ and $\epsilon > 0$. Choose $h' = \sum_{i \in I} c_i h^{\phi_i}$ such that $\|h - h'\|^2 < \epsilon/2$.

Now consider for $t \in \mathbb{R}$

$$\begin{aligned} |E(e^{itY_n(h)} - e^{itW(h)})| &\leq E|e^{itY_n(h)} - e^{itY_n(h')}| + E|e^{itY_n(h')} - e^{itW(h')}| \\ &\quad + E|e^{itW(h')} - e^{itW(h)}|. \end{aligned}$$

Using Schwartz's Inequality and the fact $|e^{ix} - 1| \leq |x|$ we get that the first

and third term of the above inequality are dominated by $t^2 E \|h - h'\|^2$ using (0.4) and (0.5). Hence by (0.3)'

$$\lim_{n \rightarrow \infty} |E e^{itY_n(h)} - E e^{itW(h)}| \leq \varepsilon/2.$$

As ε is arbitrary we get the result.

Finally, we make some observations to be used later.

$$(0.7) \quad Y_n^t(h^\phi) = \sum_{k=0}^{[nt]} n^{-k/2} \sum_{1 \leq s_1 < \dots < s_k \leq [nt]} \phi(X_{s_1}) \dots \phi(X_{s_k}) = \prod_{i=1}^{[nt]} \left(1 + \frac{\phi(X_i)}{\sqrt{n}}\right).$$

Also, $\min(t, s) \mu(A \cap A')$ is a covariance on $[0, \infty) \times \Sigma$ giving that there exists a centered Gaussian process $\underline{W}(t, A)$ with $E \underline{W}(t, A) \underline{W}(s, A') = \min(t, s) \mu(A \cap A')$. Let for $T < \infty$

$$H_T = \{(h_0, h_1, \dots) \in H : \sum_{k=0}^{\infty} T^k \frac{\|h_k\|^2}{k!} < \infty\}.$$

1. Invariance Principle: Let $D[0, T]$, $(T \leq \infty)$ be the space of right continuous functions on $[0, T]$ ($[0, \infty)$) with left limits at each $t \leq T$. The space $D[0, T]$ is endowed with Skorohod topology [1]. The topology in $D[0, \infty)$ is the one described in Whitt [4]. We note that

$X_{[nt]} = \sum_{i=1}^{[nt]} \left(\frac{\phi^2(X_i) - E\phi^2}{n} \right)$ has stationary independent increments. So for $\varepsilon > 0$

$$P\left(\sup_{0 \leq t \leq T} |X_{[nt]}| > \varepsilon\right) \leq C \cdot P(|X_{[nT]}| \geq \varepsilon) \rightarrow 0$$

by the weak law of large numbers. Using this, the arguments preceding Lemma 0.3, invariance principle and Cramér-Wold device we get the following analogue of Lemma 0.3.

Lemma 1.1: $(Y_n^t(h^{\phi_1}), \dots, Y_n^t(h^{\phi_k})) \xrightarrow{\mathcal{D}_{k,T}} (\exp(I_1^t(\phi_j) - \frac{1}{2}t\|\phi_j\|^2), j=1, \dots, k)$

where $I^t(\phi_j) = \iint 1_{(0,t]}(u) \phi_j(x) W_k(du, dx)$. Here $\xrightarrow{D_{k,T}}$ denotes convergence in $D^k[0,T]$ with respect to product topology.

We note that $W(t,A)$ is a Brownian motion for each $A \in \Sigma$. Thus we can choose $I^t(\phi)$ continuous for each ϕ and a martingale in t as $I^t(\phi) = \int \phi(x) W(t, dx)$. We get for $\{c_1, \dots, c_k\} \subseteq \mathbb{R}$, (k finite),

$$Y^t\left(\sum_{j=1}^k c_j h^{\phi_j}\right) \rightarrow \sum_{j=1}^k c_j \exp(I^t(\phi_j) - \frac{1}{2} t \|\phi_j\|^2).$$

Let $\phi \in L_0^2(X, \Sigma, \mu)$, $\|\phi\| = 1$, and denote

$$(\phi^k)^t = \phi(x_1) \dots \phi(x_k) 1_{(0,t]}(u_1) \dots 1_{(0,t]}(u_k).$$

Define $I_k(\phi^k)^t = k! H_k(t, I(\phi))$ where H_k is Hermite polynomial, i.e.

$\sum_{k=0}^{\infty} \gamma^k H_k(t, x) = \exp(\gamma x - \frac{1}{2} \gamma^2 t)$. For $\phi \in L_0^2(X, \Sigma, \mu)$, $\|\phi\| = 1$, we define for $(h^\phi)^t = (1, \phi^t, (\phi^2)^t, \dots)$,

$$W(h^\phi)^t = \sum_{k=0}^{\infty} \frac{I_k(\phi^k)^t}{k!},$$

and extend it linearly to $(\sum c_j h^{\phi_j})^t$. It is a martingale. Let $h \in H_T$ $\{h(n)\}$ a sequence in $\text{sp}\{(h^\phi)^t, \phi \text{ in CONS in } L_0^2(X, \Sigma, \mu)\} \subseteq H_T$, then

$$\begin{aligned} P(\sup_{t \leq T} |W^t(h(n)) - h(m)| \geq \varepsilon) &\leq E |W^T(h(m)) - h(n)|^2 \\ &= \sum_{k=0}^{\infty} T^k \frac{\|h_k(m) - h_k(n)\|^2}{k!} \end{aligned}$$

using Doob's inequality and argument as in (0.5). Define for $h \in H^t$,

$W^t(h) = -\lim W^t(h_n)$ where the limit is uniform on compact for $h_n \rightarrow h$. Then

$W^t(h)$ is right continuous martingale and has the same distribution as

$\sum_k I_k^t(h_k)/k!$. Now we derive the main theorem of [3].

Theorem 1.2: $Y_n^t(h) \xrightarrow{D} W^t(h)$ in $D[0,T]$ for $h \in H^T$ for each $T < \infty$.

Proof: Let $h \in H$ and $\varepsilon > 0$, choose $h'_k \in \text{sp}\{h^\phi : \phi \in L^2_0(X, \Sigma, \nu)\} \ni h_k \rightarrow h$. Now define

$$X_{nk}^\bullet = Y_n^\bullet(h'_k), Z_n^\bullet = Y_n^\bullet(h), X_k^\bullet = W^\bullet(h'_k) \text{ and } X = W^\bullet(h).$$

Then $X_{n,k}^\bullet \xrightarrow{\mathcal{D}} X_k^\bullet$ as $n \rightarrow \infty$ in $D[0, T]$ for each $T < \infty$ by Lemma 1.1. Also $X_k^\bullet \xrightarrow{\mathcal{D}} X$ as $n \rightarrow \infty$ in $D[0, T]$ for each $T < \infty$. In addition,

$$P\left(\sup_{0 \leq t \leq T} |X_{nk}^\bullet - Z_n^\bullet| \geq \varepsilon\right) \leq E|Y_n^T(h - h'_k)|^2 \leq T \|h - h'_k\|^2$$

giving $\lim_{k \rightarrow \infty} \lim_n P(\rho(X_{nk}^\bullet, Z_n^\bullet) \geq \varepsilon) \rightarrow 0$ with ρ being the Skorohod metric on $D[0, T]$. This implies by ([1], Thm 4.2, p. 25) that $Z_n^\bullet \xrightarrow{\mathcal{D}} W^\bullet(h)$ in $D[0, T]$ ($T < \infty$) giving the result.

Remark: In the above arguments we may use an interpolated version of $Y_n^t(h)$ from the beginning and use appropriate version of Donsker's Invariance Principle to conclude above convergence occurs in $D[0, T]$ in sup norm giving $W^t(h)$ continuous.

References

- [1] Billingsley, P. (1968) Convergence of probability measures, Wiley, New York.
- [2] Dynkin, E.B. and Mandelbaum, A. (1983) Symmetric statistics, Poisson point processes and multiple Wiener integrals, Ann. Statist. 11 739-745.
- [3] Mandelbaum, A. and Taqqu, M.S. (1984) Invariance principles for symmetric statistics. Ann. Statist. 12 483-496.
- [4] Whitt, W. (1980) Some useful functions for functional limit theorems, Math. of Op. Research 5 67-85.

END

4-~~2~~-87

DTIC